


Human Papillomavirus
vaccine HPV
(jag)



Easy Read Booklet

What is the vaccine (jag)


It is a liquid given to you in a jag.




The doctor or nurse will give you the jag.



Why have the HPV jag



The HPV virus is very common and you catch it by having sex.



This jag helps to prevent against cervical cancer which can grow in your cervix which is inside your vagina.




It doesn't stop all infections, so you will still need a cervical smear when 25 years of age.



If you want to get the jags you need to sign a form to say you will have it this is called a consent form.

Your HPV vaccine (jag)

If under 15 years



You will need **2** jags. Your school will tell you when you are to have your jags.

After 1st jag



2nd -6 months later.




Mark your appointments in your calendar.

If over 15 years of age



You will need **3** jags.

After 1st jag



2nd -1 month after the 1st jag.




3rd -3 months after 2nd jag.



Mark your appointments in your calendar.

Side Effects

After jag



Your arm may go a bit red, feel sore.



You may feel tired.



You may get an upset stomach or have a headache.



If you start to feel ill go to your doctor.



For more information
www.immunisationscotland.org.uk/hpv



e-mail:-
[nhs.healthscotland-
publication@nhs.net](mailto:nhs.healthscotland-publication@nhs.net)



Or call NHS inform 0800 22 44 88

Most of the photos used in this book are Photosymbols.